

## Assessment of Wheat and Quality Protein Maize Quality

Rajat Avsthi<sup>1</sup> and Kushanak Rawat<sup>2</sup>

<sup>1</sup>Department of Pharmacy, DIT University, Dehradun, INDIA.

<sup>2</sup>Department of Pharmacy, DIT University, Dehradun, INDIA.

<sup>2</sup>Corresponding Author: rawat.kush23@yahoo.com

Date of Submission: 20-07-2022

Date of Acceptance: 10-08-2022

Date of Publication: 31-08-2022

### ABSTRACT

A study was conducted to prepare protein enriched biscuits which could be used as a protein supplemented snack food. The study evaluated the effect of substituting soy-flour 20%, 15% and 10% in biscuit production. Biscuits were enriched with soy-flour T<sub>0</sub> (control with no soy flour), T<sub>1</sub> (20 %), T<sub>2</sub> (15 %) and T<sub>3</sub> (10 %). Sensory, nutrient estimation and Microbial qualities of biscuits were evaluated. Organoleptic test of biscuits showed that with regard to flavour and taste, body and texture, colour and appearance and overall acceptability, sensory characteristics of T<sub>3</sub> were found to be the best. The other treatments T<sub>0</sub>, T<sub>1</sub> and T<sub>2</sub> were also found acceptable. In nutrient estimation of biscuits, moisture content was found maximum in treatment T<sub>1</sub>, protein content was maximum in treatment T<sub>1</sub>, fat content was highest in treatment T<sub>1</sub>, ash content was highest in treatment T<sub>1</sub>, carbohydrate was found maximum in treatment T<sub>3</sub>, calorie content was maximum in T<sub>1</sub>, iron content was maximum in T<sub>2</sub> and calcium content was maximum in treatment T<sub>1</sub>. Best treatment was T<sub>1</sub> among T<sub>0</sub>, T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub>. The soy flour added product-biscuit has good shelf life. The product is ready -to-eat so don't require much time for preparation. It is a compact source of energy and nutrients including protein, fat, Iron, calcium etc. It can be concluded that Sensory scores indicated high acceptability for treated biscuit samples. Protein and fat contents of biscuits increased with increasing soy fortifications. Substitution with soy-flour did not adversely affect qualities.

**Keywords-** Soy flour, fortification, malnutrition, biscuits, organoleptic characteristics

## I. INTRODUCTION

Protein malnutrition is a serious problem in India due to cereal based dietary pattern. Therefore, various preparations based on cereal-pulse combination are of paramount importance to improve the protein quality of Indian diet. FAO (Food and Agricultural Organization) suggested that to meet the recommended dietary allowances of infants, preschool children, adolescent girls, pregnant and lactating women, low-cost supplementary foods could be processed domestically by simple, inexpensive processing technology. Some 49percent of the 10.7 million deaths among under five children each year in the developing world is associated with malnutrition. Nearly 30 percent of humanity is currently suffering from one or more of the multiple forms of malnutrition. Currently, an estimated 149.6 million children under five years of age i.e., 26.7 percent of the world's children in this age group are still malnourished when measured in terms of weight for age.

The requirement of supplementary foods is increased to reduced malnutrition. Supplementary foods should be such, if taken in small quantity, could provide necessary amount of nutrients. They should be made in form of ready-to-eat snacks, drinks etc. so that they could be taken without any inhibition. But people are still not much aware of their importance. They are just taken as conventional dishes involve either too much time or drudgery. Moreover, the foods bought from hotels and stalls are rarely of high nutritive value. Hence, the need arises for time and labor-saving convenience foods for the modern man. Soybean (*Glycine max*) the miracle bean is the base of healthy life. The nutritive virtues of soybean were intuitively realized in the orient over 4000 years ago. Where it is used in multiple foods. At the turn of last century, the west realized these virtues of soybean but the taste of foods and beverages made from soybeans were unacceptable to western palate that grossly limited their food use. More recently, soybean use is gaining acceptance in the form of textured vegetable protein (TVP), popularly sold as soy vadi and soy nuggets (American Soybean Association, 2004). Hundred grams of soybean contains 8.1 g of moisture, 43.2 g of protein, 19.5 g of fat, 20.9g of carbohydrates, 240 mg of calcium, 10.4 mg of iron and 690 mg of phosphorus (Gopalan et al., 1999). Soybean is a versatile plant food that provides high quality protein but only minimal saturated fat. However, recent excitement has focused on soy foods as a rich and essentially unique dietary source of isoflavones and phytoestrogens. Soy protein also directly lowers serum cholesterol levels (Messina, 2004). Soybean is a rich source of almost all the essential nutrients. It is commercially an oil seed but is deficient in vegetable oil. However, most of the oilseed contain 40-50 percent oil but

soybean contains about 18 percent of oil (American Soybean Association, 2004). The original impetus for research into soybean is to benefit for its high protein content i.e. about 40 percent in bean. Owing to these qualities, soybean has long been used in supplementary foods. Soybean contains isoflavones, which are said to have potential anticancer effects. It contains two primary isoflavones called as Genistein and Diadzein and a minor one called as Glycitein. They retard bone loss in premenstrual and postmenstrual women, soluble fibre in soy foods control blood sugar. Soyfoods are quite important to us as they reduce the risk of heart disease. It reduces menopausal symptoms and bone deterioration. Regular soyfood consumption delays the process of aging. Soyfood every day is protective against many types of cancers, especially breast and prostate. Also, soyfood given to children daily improves mental and physical abilities, memory power and haemoglobin levels. (American Soybean Association, 2004). Soybean was shown to be extremely rich in nearly all the essential amino acids needed by man (Cook and Briggs, 1977).

Biscuits constitute major component of human snacks in most part of the world. It is an unleavened crisp, sweet pastry made from wheat flour, shortening (hydrogenated fat) and sugar, and is usually made light by the addition of baking powder (a mixture of sodium carbonate, sodium bi phosphate and cereal flour) (O'Brien et al., 2003).

Biscuits have been man's food since a long time. It is a processed convenience food ever produced and is most widely acceptable. It is one of the few universal staples, which is complete in it and requires no additional preparation. Thus, for many, biscuit becomes an important source of high molecular carbohydrates, vegetable proteins and some vitamins and minerals. But it is important to know that, as compared to animal proteins refined wheat flour is deficient in certain essential amino acids, thus has a low nutritional value. The nutritional value of biscuit can be enhanced by fortification and supplementation with a wide variety of protein, vitamin and mineral sources. This work was designed to economically complement and fortify wheat flour and cassava with fermented or blanched soy flours for biscuit production.

**Table A: Method of preparation of Biscuits: Control and experimental samples**

Control	Experimental
20g sugar was powdered and mixed with 20 ml of fat (refined oil) for about 10 minutes.	20g sugar was powdered and mixed with 20 ml of fat (refined oil) for about 10 minutes.
50g refined wheat flour, 15 ml milk, 1/4tsp baking powder, 1/8tsp salt and 5-6 drops of vanilla essence was added and kneaded to stiff dough.	50g refined wheat flour, 15 ml milk, 1/4tsp baking powder, 1/8 tsp salt, 5-6 drops of vanilla essence, 20 percent, 15 percent and 10 percent soybean ( <i>Glycine max</i> ) flour was added to it and kneaded to stiff dough.
Dough was divided in small parts and moulded in desired shapes.	Dough was divided in small parts and moulded in desired shapes.
Tray was greased and biscuits were placed on it.	Tray was greased and biscuits were placed on it.
Biscuits were baked in pre heated oven at 140°C for 10 min.	Biscuits were baked in pre heated oven at 140°C for 10 min.

## II. MATERIALS AND METHODS

The study was carried out in the Foods and Nutrition laboratory of School of Home Science Sam Higginbotom Institute of Agriculture, Technology & Sciences, Formerly (Allahabad Agricultural Institute Deemed University).

### Sensory Analysis

Organoleptic test of the products was done by 9-point hedonic scale scorecard, especially prepared for the purpose. A 10-member trained panelists from staff and students of Halina School of Home Science Sam Higginbotom Institute of Agriculture, Technology & Sciences, Formerly (Allahabad Agricultural Institute Deemed University). Each attribute was scored based on its intensity scaled on a 9-point hedonic scale (1=disliked extremely, 2=disliked very much, 3=disliked moderately, 4=disliked slightly, 5=neither liked nor disliked, 6=liked slightly, 7 = liked moderately, 8 = liked very much, 9 = liked very extremely) for colour, flavor, mouth feel and texture.

### Assays

Biscuits were prepared using different levels of soy flour T<sub>1</sub> (20 %), T<sub>2</sub> (15%), T<sub>3</sub> (10 %) and T<sub>0</sub> was control without addition of soyflour. Analysis of products was done by AOAC methods (AOAC, 2000). Carbohydrate was calculated by difference method; energy was calculated from fat, protein incineration of sample, moisture was determined by heating ether extract method, calcium was determined titrimetrically by titration with potassium permanganate, iron was determined by spectrophotometer method. Soy flour added biscuits were analysed for total bacterial and total yeast and mold count after 15 days.

### *Statistical Analysis*

All biscuit sample data were analyzed statistically. ANOVA was used for determining the significance/ non-significance of results. SPSS 11.5 version was used to analyze the data. Results are expressed as mean & CD.

## **III. RESULTS**

### *Sensory Evaluation*

In the present study, sensory scores of biscuits enriched with 20 %, 15 % and 10 % soyflour showed that with regard to flavour and taste, body and texture, colour and appearance and overall acceptability, sensory characteristics of T<sub>3</sub> (10%) were found to be the best. Other treatments T<sub>0</sub>, T<sub>1</sub> (20%), T<sub>2</sub> (15) and T<sub>3</sub> (10%) were also found acceptable (Fig. 1.1).

### *Nutrient Estimation*

Treatment T<sub>1</sub> with addition of soy-flour had maximum protein content followed by T<sub>2</sub> and T<sub>3</sub> at 20 %, 10 and 15 % level (Table, Fig.1.2). Treatment T<sub>1</sub> with addition of soy-flour had maximum protein fat content followed by T<sub>2</sub> and T<sub>3</sub> at 20 %, 15 % and 10 % level (Table, Fig.1.3). In the present study, treatment T<sub>3</sub> with addition of soy-flour had maximum protein carbohydrate followed by T<sub>2</sub> and T<sub>1</sub> at 10 %, 15 % and 20 % level (Table, Fig.1.4). In the present study, treatment T<sub>3</sub> with addition of soy-flour had maximum iron content followed by T<sub>2</sub> and T<sub>1</sub> at 10 %, 15% and 20% level (Table, Fig.1.5). In the present study, treatment T<sub>1</sub> with addition of soy-flour had maximum calcium content followed by T<sub>2</sub> and T<sub>1</sub> at 20 %, 15 % and 10% level (Table, Fig.1.6).

## **IV. DISCUSSION**

Taste is the primary factor which determines the acceptability of any product, which has the highest impact as far as market success of product, is concerned. In the present study, overall acceptance was best for T<sub>3</sub> (Fig. 1.1). In a previous study it was observed that the mean scores for all the assessed organoleptic characteristics decreased with increase in the soybean flour. Organoleptic evaluation indicated that there were no significant ( $p < 0.05$ ) differences between the control treatment and 10% soy flour supplemented biscuits in the organoleptic attributes of taste, texture and flavour but differences were significant in colour and overall acceptability. From the overall acceptance rating, 10% soybean flour incorporated biscuit obtained the highest preference compared to other combinations (Banureka and Mahendran, 2009). In another study, the colour, flavour, mouth feel and overall acceptability of biscuits were not significantly affected by the 20%, 5%, 10% or 15% soy flour replacements for wheat flour. However, colour scores increased slightly while flavour and mouth-feel decreased slightly as the substitutes increased (Ugwuona, 2009). In another study it was observed that overall acceptability was maximum for 70% soy fortified biscuit (Kumar et al., 2010).

Soybean is a protein rich oil seed, which is presently number one edible oil source globally. Soybean is rich in polyunsaturated fats, including the two essential fatty acids, linoleic and linolenic, that are not produced in the body. Linoleic and linolenic acids aid the body's absorption of vital nutrients and are required for human health (Hegstad, 2008). In the present study, highest protein and fat content was found in T<sub>1</sub> followed by T<sub>2</sub> and T<sub>3</sub> (Table, Fig.1.2). Protein, fat and energy (calorie) value of soy flour supplemented biscuits increased with progressive increase in proportion of soy flour and 10% soy flour added biscuits obtained values of 9.9%, 20% and 453.58 kcal/g respectively, while lowest values of 5%, 14.5% and 417.36 kcal recorded for the wheat flour biscuits. The moisture and ash were decreased with corresponding increase in the percentage of soy flour (Banureka and Mahendran, 2009). In a previous study, protein and fat contents of biscuits increased with increasing soy fortifications (Ugwuona, 2009). In a study, two types of wheat flour and quality protein maize (QPM) based biscuits with and without processed defatted maize germ cake (PDMGC) supplementation were prepared and compared with wheat flour based biscuits as standard. Both protein quality and quantity improved in wheat flour + QPM based biscuits with supplementation if PDMGC at the cost of slight decrease in IVPD (Gupta and Singh, 2005).

In another study, significant decrease was counteracted by supplementing with 10% soy flour which resulted in a significant increase in the protein content. Therefore, biscuit with comparable nutrient content to 100% wheat flour could be produced with wheat composite flour up to substitution level of a minimum of 10% soy flour. This means that biscuit from wheat and soy flour composite will be acceptable in terms of colour, aroma and overall acceptability (Oluwamukomi et al., 2011).

## **V. CONCLUSION**

The finding of this re-search revealed that, the biscuits produced with soy flour substitution, upto 25%, were nutritionally superior to that of the whole wheat flour biscuits. To obtain biscuits of high nutritional and organoleptic qualities, wheat flour could be substituted with 10% of soy flour. Organoleptically, treatment T<sub>3</sub> (10%) was found to be the best and in biochemical analysis, treatment T<sub>1</sub> (20%) was found to be the best. It might be because highest percent of

soybean was added to T<sub>1</sub> (20%) and soybean has high protein, fat, iron, calcium content. As biscuit consumption in whole world is very high, soy fortified biscuits will help in increasing intake of protein, fat and calories.

### REFERENCES

- [1] American Soybean Association, 2004. United Soybean Board. "Soy and Health". A review of the 3<sup>rd</sup> International Symposium on the role of soy in preventing and treating chronic disease, October 31, 1999 in Washington DC, USA.
- [2] AOAC, 2000. Official methods of analysis of Association of Official Analytical Chemistry International, 17th Ed, Geithersburg
- Banureka V.D. and Mahendran T., 2009. Formulation of Whea-Soybean Biscuits and Their Quality Characteristics. *Tropical Agricultural Research & Extension*, 12(2): 62-66.
- [3] Cook F. and Brigg G.M., 1977. Nutritive value of eggs. In "Egg Science and Technology", WJ, Sladelman and PJ Cotteril (eds) 2nd ed. Avi Publ. Co. Inc., Connecticut: 92-108.
- [4] Gopalan C., Rama Shastri, B.U. and Balasubramaniam, S.C., 1999. Nutritive value of Indian Foods, NIN, ICMR, Hyderabad: 48.
- [5] Gupta H.O. and Singh N.N., 2005. Preparation of wheat and quality protein maize based biscuits and their storage, protein quality and sensory evaluation. *J. Food Sci. Tech.*, 42(1): 43-46.
- [6] Hegstad H.G., 2008. Nutritional and Health benefits of Soybean. Soy Protein Quality Evaluation Report, Food and Agriculture Organization of the United Nations, Food and Nutrition, Rome, Italy, Paper No. 71.
- [7] Kuar S., Rekha and Sinha L.K., 2010. Evaluation of quality characteristics of soy based millet biscuits. *Advances in Applied Science Research* ,1(3): 187-196. Messina M., 2004. Defining Isolavones. *Soy and Health*, www.asaasc.com> viewed 2011.
- [8] O'Brien C.N., Champmemb O., Nexille D.P., Kengh M.K., and Arendt E.K., 2003. Effect of varying micro-encapsulation process on the functionality of hydrogenated vegetable fat in short dough biscuit. *Food Research International*, 36: 215-221.
- [9] Oluwamukomi M.O., Oluwalana I.B., and Akinbowale O.F., 2011. Physicochemical and sensory properties of wheat cassava composite biscuit enriched with soy flour. *African Journal of Food Science*, 5 (2): 50 - 56.
- [10] Ugwuona F.U., 2009. Chemical And Sensory Evaluation of Soy-Fortified Cassava wheat Biscuit. *Journal of Tropical Agriculture, Food, Environment and Extension*, 8 (1): 55-59.