

Chemical Processes and the Effects on the Environment of Converting Heavy Hydrocarbons to Gasoline

Abdulsamea Mohammed Elhadi Ali Hussen¹ and Mohamed Abusa Gaza²

¹Department of Preschools, Faculty of Education, Zintan University, Derj, Ghadames, LIBYA

²Department of Biotechnology, Faculty of Health Sciences, Aljafara University, Alzahram, LIBYA

¹Corresponding author: abdulsamea.hussen@uoz.edu.ly



www.sjmars.com || Vol. 1 No. 1 (2022): February Issue

Date of Submission: 30-01-2022

Date of Acceptance: 25-02-2022

Date of Publication: 28-02-2022

ABSTRACT

The conversion of heavy hydrocarbons to gasoline is a crucial process in the petroleum industry, utilizing various refining techniques such as catalytic cracking, hydrocracking, thermal cracking, and coking. This study looks at the environmental effects of the chemical processes that turn heavy hydrocarbon molecules into lighter products like gasoline. After a thorough literature review, this study uses qualitative methodology to examine the impact of the chemical process that converts heavy hydrocarbons to gasoline on the environment. This study's systematic approach to literature selection prioritizes peer-reviewed sources and respected industry publications to ensure the validity and trustworthiness of the results. According to the results, converting heavy hydrocarbons into gasoline is one of the most essential operations in the petroleum industry as it ensures fuel supply for transportation and industry. Coking, hydrocracking, thermal cracking, and catalytic cracking are some chemical processes that pose significant environmental risks. These activities cause climate change and global warming because Carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) are among the greenhouse gases released. These efforts result in air pollution, elements that cause smog, acid rain, and other adverse health effects, such as sulfur oxides (SO₃), nitrogen oxides (NO₃), and particulate matter.

Keywords- Chemical, Processes, Environment, Heavy Hydrocarbons, Gasoline.

I. INTRODUCTION

Turning heavy hydrocarbons into gasoline is one of the most essential steps in the petroleum refining process. Many chemical reactions, including as catalytic cracking, hydrocracking, coking, and visbreaking, are principally responsible for this change.¹ Zeolite-based catalysts are currently being researched for hydrocarbon synthesis (Figure 1). This technique divides large, complex hydrocarbon molecules into smaller, more valuable products such as diesel, gasoline, and other light fractions.




Figure 1. Catalysts for the conversion of CO₂ into hydrocarbon fuels, both zeolite and non-zeolite²

The refining method has a major impact on the environment despite being necessary to meet the world's energy needs. This process's pollutants, waste products, and energy use cause air pollution, greenhouse gas emissions, and possible ecological damage. Understanding the environmental impacts of this chemical process is necessary to develop more environmentally friendly refining methods and reduce adverse environmental impacts.² Heavy hydrocarbons can be converted into gasoline primarily through catalytic cracking.

This procedure uses a catalyst, often a zeolite-based material to aid in the controlled breakdown of heavy molecules. In modern oil refineries, fluid catalytic cracking (FCC) is the most widely used technique for efficiently turning heavy oil into gasoline and other valuable products. However, significant amounts of carbon dioxide (CO₂), sulfur oxides (SO_x), and nitrogen oxides (NO_x) are discharged by FCC units into the environment. These pollutants contribute to global warming, acid rain, and respiratory issues in people. To prevent environmental damage, hazardous waste resulting from the use of catalysts containing heavy metals such as nickel and vanadium must be recycled or disposed of properly.³

Refining heavy hydrocarbons also involves a critical process called hydrocracking, which breaks down heavy molecules by applying high pressure and temperature to hydrogen and a catalyst. Compared to catalytic cracking, hydrocracking produces fewer unwanted byproducts, such as sulfur compounds, making it a cleaner process. However, a significant quantity of hydrogen is needed for this process, and steam methane reforming (SMR) is usually used to create hydrogen from natural gas. SMR adds to the total carbon footprint of hydrocracking as it is a very energy-intensive process that emits much CO₂. Additionally, hydrocracking facilities produce wastewater containing sulfides, ammonia, and other impurities that must be treated appropriately before discharge to avoid contaminating water supplies.⁴

Another popular technique for handling heavy hydrocarbons is coking. High-temperature thermal cracking creates lighter fractions while solid petroleum coke is left behind. Liquid and delayed coke are the two main coking processes used in oil refineries. In addition to producing large quantities of petroleum coke, a carbon-rich material often used as a fuel in industrial settings, cokers efficiently convert waste oil into usable fuel. Heavy metals and large amounts of CO₂ are released into the atmosphere when petroleum coke is burned, accelerating climate change and increasing air pollution.⁵

In addition, the health of nearby residents is affected by volatile organic compounds (VOCs) and particulate matter produced by coking units. The purpose of visbreaking, a milder thermal cracking method, is to reduce the viscosity of the heavy residue and create a lighter distillate. However, visbreaking produces significant amounts of gas and liquid byproducts, although not as severe as coking or catalytic cracking. The effluent from visbreaking contains hydrocarbons, hydrogen sulfide (H₂S), and other contaminants that must be cleaned up to meet environmental regulations. The wastewater from visbreaking facilities may contain polycyclic aromatic hydrocarbons (PAHs), which are known carcinogens and can affect water bodies if not adequately controlled.⁶

The environmental impact of converting heavy hydrocarbons into gasoline goes beyond refinery emissions and waste products. This refining process primarily uses crude oil as its raw material, and the extraction and transportation of this fuel significantly negatively impacts the environment.⁷ Oil drilling operations usually have detrimental impacts on land and marine habitats, including habitat loss, contamination of water and soil, and oil spills. Transporting crude oil through pipelines and tankers increases the likelihood of leaks and spills, exacerbating environmental problems. Examining how chemical reactions that transform heavy hydrocarbon molecules into lighter compounds, like gasoline, affect the environment is the goal of this study.

II. LITERATURE REVIEW

2.1. Gasoline

Gasoline is widely used in internal combustion engines that power cars, motorcycles, light aircraft, ships, and other machines. Gasoline is a refined petroleum-based fuel. Gasoline's complex hydrocarbon composition makes it highly combustible and energy dense.⁸ Typically, gasoline contains compounds with carbon chains ranging from C₄ to C₁₂. Several refining techniques are used to manufacture gasoline, including fractional distillation, catalytic cracking, hydrocracking, and reforming. All of these techniques convert heavier hydrocarbons into lighter, more combustible fractions. Paraffins, naphthenes, and aromatic hydrocarbons make up the majority of gasoline's chemical makeup, albeit this varies based on the source of the crude oil, refining methods, and regulatory requirements (Figure 2).⁹




Figure 2. Schematic of Refinery Products.⁹

The octane rating of gasoline is one of its most essential properties; it indicates how resistant the fuel is to engine knock, a condition caused by premature combustion in the engine cylinders. Premium gasoline and other high-octane fuels are recommended for high-performance or high-stress engines because of their smoother combustion characteristics. Additives are often used to improve the economy of gasoline and its environmental performance. These additives include oxygenates such as ethanol to improve combustion and reduce emissions, detergents to remove engine deposits, and antioxidants to stop fuel breakdown.¹⁰

Although gasoline is essential to modern industry and transportation, the impact on the environment is very detrimental. Nitrogen oxides (NO₂), carbon dioxide (CO₂), and volatile organic compounds (VOCs) are produced when fuel is used, all of which exacerbate air pollution, smog, and climate change. In addition, gasoline can leak or spill during production, transportation, or storage, contaminating surrounding soil and water. As concerns about environmental sustainability and dependence on fossil fuels grow, efforts to develop alternative fuels, improve fuel economy, and move to electric and hybrid vehicles are influencing the future of gasoline consumption and global energy use¹¹

2.2. Heavy Hydrocarbons

High molecular weight and lengthy carbon chains are characteristics of heavy hydrocarbons; most atoms in these complex organic compounds are hydrogen and carbon. These hydrocarbons are often found in natural fossil fuels such as bitumen and crude oil. These include heavy crude oil, asphalt, residual fuel oil, and waxes, among other substances distinguished by their high viscosity, low volatility, and resistance to evaporation. Heavy hydrocarbons are converted into more valuable and profitable goods such as diesel, gasoline, and petrochemicals, due to their complex structure, a large amount of processing and refining is required.¹²

Coking, visbreaking, hydrocracking, thermal, and catalytic cracking are essential to refining heavy hydrocarbons. This process breaks down large hydrocarbon molecules into lighter fractions with lower boiling points, making them more suitable for transportation and combustion. However, the energy-intensive refining of heavy hydrocarbons often results in significant carbon dioxide (CO₂), sulfur oxides (SO_x), and nitrogen oxides (NO_x) emissions, which worsen air pollution and fuel climate change. To further lessen their detrimental impacts on the environment, solid waste byproducts from the processing of these hydrocarbons, such as petroleum coke and heavy metal wastes, must be recycled or disposed of correctly (Figure 3).¹³




Figure 3. Developments of adsorbents for the separation and purification of hydrocarbons¹³

Many industries rely on heavy hydrocarbons to produce chemicals, lubricants, and gasoline. Heavy hydrocarbons are widely used in electricity generation, road construction, and naval fuels due to their extreme durability and high energy content, which offer substantial industrial advantages. However, its extraction, transportation, and processing incur environmental costs, including resource depletion, greenhouse gas emissions, and water and soil contamination due to spills. As global energy demand changes, efforts are being made to reduce dependence on heavy hydrocarbons and increase the sustainability of their use in modern industry through innovations in refining technology and other energy sources.¹⁴

2.3. Chemical Processes

In many industrial applications, a chemical process is a series of chemical reactions and physical changes that transform raw materials into desired products. Through chemical reactions, separation methods, and energy transfer mechanisms, these processes change a substance's molecular structure, content, and characteristics. Chemical processes are essential in environmental engineering, manufacturing, food production, pharmaceuticals, and petroleum refining. Depending on the type of reaction and the desired end product, chemical processes can be divided into several categories, such as synthesis, cracking, oxidation-reduction, polymerization, and catalysis.¹⁵

Chemical processes in an industrial context are made as efficient, productive, and waste-free as possible. Typical examples are electrolysis, which uses electrical energy to drive a chemical process; distillation, which separates components based on their boiling points; and catalytic cracking, which converts heavy hydrocarbons into lighter fuels. Catalysts, chemicals that increase the reaction rate without being consumed, are used in many of these processes to increase efficiency and selectivity. Furthermore, chemical engineering principles including reaction kinetics, thermodynamics, and process control are used to ensure safe and efficient operations in large-scale chemical production.¹⁶

Chemical processes provide benefits, but they can pose safety and environmental risks. Many industrial operations produce emissions, hazardous wastes, and by-products that can exacerbate environmental degradation, air and water pollution, and climate change. Therefore, companies are increasingly using greener chemical processes that use renewable resources, energy-efficient techniques, and waste reduction tactics. Biocatalysis, carbon capture, and sustainable solvents are examples of advances in chemical process technology that seek to reduce environmental impacts while maintaining productivity. Reducing resource consumption, increasing industrial sustainability, and addressing global environmental issues depend on developing new, sustainable chemical processes.¹⁷

2.4. Converting Heavy Hydrocarbons to Gasoline

Large and complex hydrocarbon molecules found in tar sands, crude oil, or waste fuels must be broken down into lighter, smaller fractions suitable for use as gasoline to refine heavy hydrocarbons into gasoline fuel. The conversion of heavy hydrocarbons into more valuable and usable fuels requires complex chemical and thermal processing due to their significant molecular weight, low volatility, and high viscosity. This conversion is essential in the petroleum industry because it maximizes the production of highly desirable products such as jet fuel, diesel, and gasoline from crude oil.¹⁸

This transformation is achieved through key refining techniques, such as coking, hydrocracking, thermal cracking, and catalytic cracking. Under controlled temperature and pressure, catalytic cracking breaks large hydrocarbon chains into smaller, more volatile molecules using catalysts such as zeolites. In contrast, hydrocracking refines heavy hydrocarbons into lighter products while removing impurities such as nitrogen and sulfur using hydrogen gas and catalysts. High heat is used in thermal cracking to break molecular bonds, producing a mixture of hydrocarbons from the gasoline series and other byproducts. Another method for removing lighter hydrocarbons is coking, which produces petroleum coke, a solid carbon residue.¹⁹

This conversion procedure has an environmental impact, but is necessary to meet the world's fuel needs. In addition to producing hazardous waste and contributing to air pollution, refining and upgrading heavy hydrocarbons releases greenhouse gases such as carbon dioxide (CO₂) and methane (CH₄). Furthermore, spills and ecological damage can arise from the extraction and transportation of heavy hydrocarbons. Research into improving refining efficiency, cutting emissions, and creating alternative fuel sources to reduce reliance on heavy hydrocarbons for gasoline production is ongoing as the world moves toward greener energy options.²⁰

III. METHODOLOGY

Using qualitative methods derived from a comprehensive literature review, this study investigates the chemical reactions that convert heavy hydrocarbons into gasoline and their environmental impact. To include up-to-date information and offer an in-depth understanding of the topic, the research will thoroughly assess scientific journals, industry reports, books, and government publications. The primary refining methods, including catalytic cracking, hydrocracking, and thermal cracking, are identified in the literature review along with their effectiveness, related technical developments, and environmental impacts. The qualitative approach of this methodology allows a thorough investigation of the many viewpoints, ideas, and technical advances on the subject.²¹

Through a rigorous evaluation of the literature, this research analyzes potential sustainable alternatives, identifies knowledge gaps, and emphasizes essential obstacles in refining heavy hydrocarbons. By integrating sophisticated in situ/operando spectroscopy and microscopy characterizations with density functional theory computations, the structure–performance correlations of common catalytic materials employed for the CO₂ conversion processes have been made public. In the future, catalytic selectivity toward a single CO₂ reduction product or fraction should be further enhanced at a CO₂ conversion rate that is relevant to industry and has a significant degree of stability (Figure 4).²¹




Figure 4. Performance Correlations of Typical Catalytic Materials Used for the CO₂ Conversion

IV. RESULT AND DISCUSSION

4.1. Result

The study's findings, which were derived from a thorough literature review, emphasize the chemical reactions involved in turning heavy hydrocarbons into gasoline and the effects these reactions have on the environment. According to the study, the main refining techniques for converting heavy hydrocarbons into lighter fuel products are thermal cracking, hydrocracking, and catalytic cracking. Catalytic cracking is the most extensively utilized method because it is efficient and can increase gasoline supply while avoiding unwanted byproducts. Hydrocracking is necessary in producing cleaner fuels despite being more expensive, since it provides benefits including improved fuel quality and lower sulfur content. Despite its effectiveness, thermal cracking and coking produce large amounts of carbon leftovers and greenhouse gas emissions, exacerbating environmental issues.²²

The study also reveals that turning heavy hydrocarbons into gasoline is still energy-intensive and has significant environmental effects, even with improvements in refining technology. The primary pollutants emitted during refining are carbon dioxide (CO₂), nitrogen oxides (NO_x), sulfur oxides (SO_x), and volatile organic compounds (VOCs), which all contribute to air pollution and climate change. Environmental deterioration is further exacerbated by the dangers associated with the extraction and processing of heavy hydrocarbons, which include soil contamination, water pollution, and oil spills.²³

The literature also mentions current studies and technical advancements meant to lessen the effects of refining on the environment. Potential strategies for improving sustainability in gasoline manufacturing include using carbon capture technology, better catalysts, and cleaner refining techniques. To lessen dependency on fossil fuels, other energy sources including biofuels and synthetic fuels are also being investigated as possible substitutes for conventional gasoline. The study's findings highlight the necessity of ongoing innovation and more stringent environmental laws to reduce the adverse consequences of turning heavy hydrocarbons into gasoline while maintaining industrial efficiency and energy security.²⁴

4.2. Discussion

4.2.1. Chemical Processes of Converting Heavy Hydrocarbons to Gasoline

Chemical Processes of Converting Heavy Hydrocarbons to Gasoline

Large hydrocarbon molecules are broken down into smaller, more valuable parts through various chemical and physical changes during the conversion of heavy hydrocarbons into gasoline. Heavy hydrocarbons require sophisticated refining methods to improve their suitability as fuels due to their high molecular weight, low volatility, and high viscosity. Coking, hydrocracking, thermal cracking, and catalytic cracking are the primary chemical processes involved in this conversion. All these techniques are essential for maximizing fuel output, increasing efficiency, and resolving environmental issues with waste management and emissions.²⁵

Catalytic cracking, which uses a catalyst usually zeo-lites to split long-chain hydrocarbons into smaller molecules at low pressures and temperatures, is one of the most popular refining techniques. This procedure is very important for fuel refining since it maintains high octane rating while increasing gasoline yield. Dehydrogenation, isomerization, and fragmentation are some processes that hydrocarbon molecules may undergo at the catalyst's active site. Since catalytic cracking uses less energy and works effectively at relatively lower temperatures, it is preferred over thermal cracking. Air pollution and other environmental problems are caused by emissions of carbon monoxide (CO), carbon dioxide (CO₂), and other volatile organic compounds (VOCs).²⁶

Hydrocracking is another vital step in converting heavy hydrocarbons into gasoline, which breaks down large hydrocarbon molecules into more minor hydrogen-saturated compounds using hydrogen and a catalyst. Unlike catalytic cracking, hydrocracking occurs in the presence of hydrogen gas under high pressure, which helps remove metals, sulfur, and nitrogen, among other contaminants. This process produces a cleaner, more stable fuel product with less sulfur, making it more environmentally friendly. Additionally, hydrocracking produces fewer unwanted byproducts while increasing gasoline and diesel production. Despite its benefits, hydrocracking is an energy-intensive procedure that uses much hydrogen. It can increase greenhouse gas emissions unless the hydrogen is produced using low-carbon or renewable resources.²⁷

Another refining technique for converting heavy hydrocarbons into gasoline is thermal cracking. This technique uses free radical processes to break down large hydrocarbon molecules at high temperatures, usually above 450°C. Thermal cracking is more adaptable to processing a variety of feedstocks because it does not require a catalyst like catalytic methods. Steam cracking and visbreaking are the two primary forms of thermal cracking. Steam cracking produces mostly lighter hydrocarbons such as ethylene and propylene, which are essential in the petrochemical industry, while visbreaking is used to reduce the viscosity of heavier residues. However, thermal cracking is less selective than hydrocracking and catalytic cracking, and often produces large amounts of coke, a solid carbonaceous residue that reduces the efficiency of the refining unit. Furthermore, the process produces large amounts of carbon dioxide emissions, exacerbating environmental problems related to climate change.²⁸

Another method for converting heavy hydrocarbons into lighter fractions is coking. With this process, heavy residual feedstocks are heated to high temperatures without oxygen, which causes thermal breakdown and the production

of lighter hydrocarbons as a byproduct, along with petroleum coke. Liquid coke and delayed coke are the two primary forms of coke. While liquid coke is a continuous process that increases efficiency and reduces downtime, delayed coke is a batch method that allows the coke to develop in the reactor. While it increases the yield of gasoline and diesel, coking also produces a large amount of petroleum coke, which is often used as a low-quality fuel with high carbon emissions. Concerns over its environmental impact are growing, especially concerning air pollution and global warming.²⁹

Due to the large volumes of greenhouse gases and other pollutants produced during the refining process, converting heavy hydrocarbons to gasoline poses some environmental concerns. Among the primary emissions produced by this process are particulate matter, carbon dioxide (CO₂), sulfur oxides (SO₃), and nitrogen oxides (NO₃). The refining sector is under increasing pressure to adopt cleaner and more sustainable technologies due to the contribution of these pollutants to acid rain, climate change, and air quality degradation. Several refining process innovations are being investigated to reduce these environmental impacts. For example, the efficiency of catalytic and hydrocracking has increased due to the introduction of advanced catalysts with reduced energy requirements and increased selectivity. Furthermore, refining facilities are incorporating carbon capture and storage (CCS) technology to capture and retain CO₂ emissions before they are released into the environment.³⁰

Other fuel sources such as synthetic and biofuels are also being researched as possible alternatives to gasoline made from heavy hydrocarbons. Biofuels are a lower-carbon substitute for traditional gasoline from renewable resources like vegetable oils and agricultural waste. Synthetic fuels from hydrogen and carbon capture technology could also offer greener energy sources with lower emissions. Ongoing research and regulatory measures are driving the shift to more sustainable fuels, although these alternatives are not yet widely used (Figure 5).³¹




Figure 5. Flowsheet for methane-to-olefins employing SMR, DRM, and POX reforming technologies³¹

4.2.2. The Effects of the Chemical Processes of Converting Heavy Hydrocarbons to Gasoline on the Environment

One of the most essential steps in the petroleum industry is the conversion of heavy hydrocarbons into gasoline, which ensures a steady fuel supply for industrial and transportation applications. However, there are severe environmental impacts from the chemical processes involved, including coking, hydrocracking, thermal cracking, and catalytic cracking. These operations use energy, emit greenhouse gases, and create many pollutants that contaminate air, water, and land. It is essential to understand these environmental impacts to develop plans to reduce harm while maintaining energy security.³²

Greenhouse gas (GHG) emissions from refining operations are among the most pressing environmental concerns. Significant volumes of carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O) are released during the combustion of fossil fuels in refining facilities and the chemical processes that occur during cracking and hydrocracking. All of these emissions play a role in climate change and global warming. Energy-intensive procedures such as hydrocracking and thermal cracking, which require high temperatures and pressures to break down complex hydrocarbons, produce carbon dioxide primarily as a byproduct. Although emitted in lower amounts, methane and nitrous oxide have a much greater potential to cause global warming than carbon dioxide, making their release particularly concerning. Rising temperatures, severe weather patterns, and ecological disruption are consequences of climate change, which is accelerated by the buildup of these gases in the atmosphere.³³

Air pollution is another major environmental problem associated with these refining operations, in addition to GHG emissions. Pollutants such as sulfur oxides (SO_x), nitrogen oxides (NO_x), particulate matter (PM), and volatile organic compounds (VOCs) are released when heavy hydrocarbons are converted. Acid rain, which damages crops, damages infrastructure, and kills aquatic life, is a result of sulfur and nitrogen oxides. Particulate matter, a combination of minor liquid and solid particles in the air, is a significant health hazard because it can cause cardiovascular disorders, respiratory problems, and premature death. In the meantime, ground-level ozone and smog are created by volatile organic compounds emitted during the refining and storage of gasoline, which deteriorates air quality and human health.³⁴

Another critical effect of turning heavy hydrocarbons into gasoline is water contamination. These operations produce large amounts of wastewater, frequently containing heavy metals, hazardous chemicals, and oil residues. This effluent can endanger aquatic life and human populations by contaminating rivers, lakes, and groundwater supplies if it is not adequately cleaned. For instance, catalysts that may leach metals like nickel, vanadium, and arsenic into wastewater streams are necessary for hydrocracking and catalytic cracking. Furthermore, thermal pollution from the discharge of cooling water used in refining facilities can disturb aquatic habitats by changing the oxygen and temperature of the water. Crude oil and refined petroleum product spills that happen by accident worsen water contamination and harm freshwater and marine ecosystems over time.³⁵

Another environmental issue related to these chemical processes is soil pollution. High concentrations of carbon, sulfur, and heavy metals are found in the solid leftovers left over after refining, including sludge, wasted catalysts, and petroleum coke. If these residues are not properly disposed of, hazardous materials may build up in the soil, decreasing its fertility and harming microorganisms and plants. Although petroleum coke, a byproduct of coking operations, is frequently used as a low-cost fuel substitute, burning it increases emissions due to its high carbon content. Furthermore, leaks and spills are common at oil storage facilities and refinery sites, which can cause groundwater pollution and long-term soil deterioration.³⁶

Energy expenditure is another crucial element influencing the environmental effects of turning heavy hydrocarbons into gasoline. Because of the high temperatures and pressures needed for the operations, fossil fuels are used extensively to generate heat. In addition to increasing reliance on non-renewable resources and contributing to increased greenhouse gas emissions, this high energy demand exacerbates the depletion of natural reserves. Since the refining industry is one of the most significant industrial energy users, increasing energy efficiency is essential to lessening its environmental impact.³⁷

Despite these obstacles, technological developments and environmental laws propel attempts to lessen the adverse consequences of refining procedures. Cleaner catalysts have reduced emissions and minimized waste output by increasing the efficiency of hydrocracking and catalytic cracking. Some refineries use carbon capture and storage (CCS) technology to absorb and store CO₂ emissions before they are released into the sky. Furthermore, water contamination from refinery effluent is decreasing with water treatment technologies, such as chemical neutralization and improved filtration.³⁸

In addition to technological advancements, the move to alternate fuels is accelerating. Alternatives to gasoline made from heavy hydrocarbons, including hydrogen-based energy sources, biofuels, and synthetic fuels, are being investigated. By significantly lowering carbon emissions and dependency on fossil fuels, these alternatives might contribute to developing a more sustainable energy environment. Additionally, international agreements and stronger environmental regulations are pushing refineries to use more sustainable practices, such as lowering the sulfur content of fuels and increasing energy efficiency through process optimization (Figure 6).³⁹




Figure 6. The Flowchart Showing The Basic Materials And Fuel Products⁴⁰

4.2.3. Strategies to Mitigate the Negative Effects of the Chemical Processes of Converting Heavy Hydrocarbons to Gasoline on the Environment

Turning heavy hydrocarbons into gasoline is essential for supplying the world's fuel needs. Still, it also presents serious environmental problems, such as greenhouse gas (GHG) emissions, soil contamination, and air and water pollution. The petroleum sector must implement measures to lessen the adverse consequences of refining operations as worries about climate change and environmental degradation grow. Advanced refining technologies, carbon capture and storage (CCS) systems, alternative fuel research, energy efficiency enhancements, and more stringent environmental restrictions are some of these tactics. By combining these strategies, refineries may lessen their environmental impact and preserve energy security.⁴¹

Using cutting-edge refining technology that increases efficiency and lowers emissions is one of the best strategies to lessen the environmental impact of refining. For instance, advancements in hydrocracking and catalytic cracking techniques have produced high-performance catalysts that reduce sulfur oxide (SO_x) and nitrogen oxide (NO_x) emissions while increasing gasoline output. By minimizing the generation of undesirable byproducts, modern catalysts like zeolites with improved selectivity help reduce waste and enhance fuel quality. Furthermore, research is being done on converting heavy hydrocarbons into cleaner-burning syngas, which may be utilized to generate electricity with less emissions, using integrated gasification combined cycle (IGCC) technology.⁴¹

Carbon capture and storage (CCS) technology is another crucial tactic. In CCS, carbon dioxide (CO_2) emissions from refinery operations are captured and stored underground or used for industrial purposes. This method can drastically lower refineries' carbon footprint by keeping CO_2 from being emitted into the atmosphere. Several refineries are also experimenting with carbon utilization strategies to lessen further their need for goods derived from fossil fuels, such as turning collected CO_2 into synthetic fuels or chemicals. However, the high price and requirement for extensive infrastructure construction make widespread implementation of CCS difficult.⁴²

Switching to alternate fuels is another vital tactic to lessen the environmental impact of turning heavy hydrocarbons into gasoline. Using renewable resources like algae and plant-based components, biofuels like ethanol and biodiesel provide a more environmentally friendly choice. Because biofuels are biodegradable and emit fewer carbon emissions than conventional gasoline, there is less chance of environmental pollution. Furthermore, as a greener substitute, synthetic fuels made from hydrogen and captured carbon are attracting attention. The industry may lessen total emissions and its reliance on petroleum-based goods by funding the study and development of these alternative fuels.⁴³

Improving refineries' energy efficiency is also crucial to lessening their environmental impact. High temperatures and pressures are necessary for the energy-intensive process of refining heavy hydrocarbons, which significantly uses fossil fuels. Refineries can reduce the requirement for extra energy input by capturing and reusing surplus heat generated during processing by implementing waste heat recovery systems. Furthermore, using renewable energy sources for refinery operations, such as solar and wind power, can assist in reducing emissions related to energy use. To reduce their dependency on fossil fuels for heating, several refineries are already using solar thermal technology to warm crude oil.⁴⁴

Another crucial component of reducing environmental damage is addressing soil and water contamination. Large volumes of effluent with heavy metals, oil residues, and hazardous compounds are produced by refineries. Advanced wastewater treatment technologies like membrane filtration and bioremediation can help keep water sources clean by removing contaminants before disposal. Using closed-loop water recycling systems can also reduce the amount of freshwater used and wastewater produced. Refineries must follow strict waste disposal guidelines to preserve soil by preventing hazardous materials from leaking into the environment from used catalysts, sludge, and petroleum coke.⁴⁵

Ignoring how government rules and policies influence environmentally friendly refining techniques is impossible. Several nations have imposed more stringent environmental regulations to reduce air pollution, such as mandating low-sulfur fuels. Refineries are compelled to use cleaner technology and reduce their emissions by regulatory frameworks like the Paris Agreement and the International Maritime Organization's (IMO) sulfur quota. To lessen the financial burden of switching to greener methods, governments also offer incentives to refineries who engage in carbon capture and renewable energy projects.⁴⁶

Corporate sustainability activities are essential for reducing environmental impact, in addition to technical and regulatory measures. Numerous oil and gas firms are investing in research for better refining techniques and pledging to meet net-zero emissions standards. Innovative solutions may be developed more quickly when industrial players, academic institutions, and environmental groups work together. Refineries' long-term environmental gains are fueled by adopting more sustainable processes due to consumer demand for greener fuels and public awareness (Figure 7).⁴⁷




Figure 7. Production, Storage, Use, And Environmental Effects of Hydrogen⁴⁷

V. CONCLUSION

In the petroleum business, turning heavy hydrocarbons into gasoline is a crucial step that guarantees fuel supply for industrial and transportation uses. However, there are serious environmental issues with the chemical processes involved, such as coking, hydrocracking, thermal cracking, and catalytic cracking. These activities contribute to climate change and global warming by releasing greenhouse gases such as nitrous oxide (N₂O), carbon dioxide (CO₂), and methane (CH₄). They also produce particulate matter, sulfur oxides (SO_x), nitrogen oxides (NO_x), and other atmospheric pollutants that cause smog, acid rain, and other adverse health impacts. Used catalysts, petroleum coke, and wastewater discharge contaminating water and soil, worsening environmental issues. Although improvements in carbon capture and storage (CCS), alternative fuels, and refining technologies provide encouraging alternatives, more extensive steps are required to reduce these effects.

Several suggestions should be considered to mitigate the environmental effects of refining heavy hydrocarbons. Refineries should prioritize implementing cleaner and more effective refining technologies to lower emissions and waste, such as enhanced catalytic systems and integrated gasification combined cycle (IGCC) technology. Second, carbon capture and utilization tactics had to be extended to reduce greenhouse gas emissions and investigate novel approaches to turning CO₂ into valuable goods. Third, more funding for alternative fuels, such as synthetic and biofuels, can offer sustainable alternatives that lessen dependency on gasoline derived from fossil fuels. Fourth, energy efficiency enhancements, such as waste heat recovery and incorporating renewable energy sources, should be implemented to reduce the carbon footprint of refining processes. Last but not least, to guarantee adherence to sustainability requirements, boost cleaner fuel production, and stimulate research into ecologically friendly refining techniques, stronger environmental policies and industrial regulations must be implemented. Following these suggestions, the petroleum sector may ensure long-term sustainability and lessen its ecological effect while balancing fuel production and environmental responsibility.

ACKNOWLEDGEMENT

I want to express my deep gratitude to my adviser for their help and advice during this research. I also want to thank the instructors and volunteers who contributed their time and knowledge to enable this research. Finally, I want to thank my family and friends for their constant support and encouragement.

DISCLOSURE

Only those who have requested disclosure can access this area. If the experts who evaluated your work agree to sell disclosure to you, their names will be listed here. Each reviewer can choose their price because it is a public endorsement of your findings and may be utilized for many purposes.

REFERENCES

- [1] Akhtar, M. S., Ali, S., & Zaman, W. (2020). Recent Advancements in Catalysts for Petroleum Refining. *Catalysts*, 14(12), 841. <https://doi.org/10.3390/catal14120841>
- [2] Garba, M. D., Usman, M., Khan, S., Shehzad, F., Galadima, A., Ehsan, M. F., Ghanem, A. S., & Humayun, M. (2021). CO₂ towards fuels: A review of catalytic conversion of carbon dioxide to hydrocarbons. *Journal of Environmental Chemical Engineering*, 9(2), 104756. <https://doi.org/10.1016/j.jece.2020.104756>
- [3] Dobre, T., Isopencu, G. O., Bdaiwi Ahmed, S., & Deleanu, I. M. (2020). Heavy Metal Pollution and Solutions for Its Control: General Aspects with a Focus on Cobalt Removal and Recovery from Aqueous Systems. *ChemEngineering*, 8(6), 118. <https://doi.org/10.3390/chemengineering8060118>
- [4] Marouani, I., Guesmi, T., Alshammari, B. M., Alqunun, K., Alzamil, A., Alturki, M., & Hadj Abdallah, H. (2020). Integration of Renewable-Energy-Based Green Hydrogen into the Energy Future. *Processes*, 11(9), 2685. <https://doi.org/10.3390/pr11092685>
- [5] Yasemi, S., Khalili, Y., Sanati, A., & Bagheri, M. (2022). Carbon Capture and Storage: Application in the Oil and Gas Industry. *Sustainability*, 15(19), 14486. <https://doi.org/10.3390/su151914486>
- [6] Haneef, T., Mustafa, M. R., Wan Yusof, K., Isa, M. H., Bashir, M. J., Ahmad, M., & Zafar, M. (2020). Removal of Polycyclic Aromatic Hydrocarbons (PAHs) from Produced Water by Ferrate (VI) Oxidation. *Water*, 12(11), 3132. <https://doi.org/10.3390/w12113132>
- [7] Geethani, M., & Kulatunga, A. (2023). Life Cycle Assessment of the Gasoline Supply Chain in Sri Lanka. *Sustainability*, 16(24), 10933. <https://doi.org/10.3390/su162410933>
- [8] Yue, Z., & Liu, H. (2020). Advanced Research on Internal Combustion Engines and Engine Fuels. *Energies*, 16(16), 5940. <https://doi.org/10.3390/en16165940>

- [9] Saha, S., Lakhe, P., Mason, M.J. et al. (2021). Sustainable production of graphene from petroleum coke using electrochemical exfoliation. *npj 2D Mater Appl*, 5, 75. <https://doi.org/10.1038/s41699-021-00255-8>
- [10] Wu, G., Ge, J. C., & Choi, N. J. (2020). Effect of Ethanol Additives on Combustion and Emissions of a Diesel Engine Fueled by Palm Oil Biodiesel at Idling Speed. *Energies*, 14(5), 1428. <https://doi.org/10.3390/en14051428>
- [11] Mohammad, M. J., Krishna, P. V., Lamma, O. A., & Khan, S. (2015). Analysis of water quality using Limnological studies of Wyra reservoir, Khammam district, Telangana, India. *Int. J. Curr. Microbiol. App. Sci*, 4(2), 880-895.
- [12] Ahmednooh, M., & Menezes, B. (2012). Blend Scheduling Solutions in Petroleum Refineries towards Automated Decision-Making in Industrial-like Blend-Shops. *Processes*, 12(3), 561. <https://doi.org/10.3390/pr12030561>
- [13] Wu, Y., & Weckhuysen, B. M. (2021). Separation and Purification of Hydrocarbons with Porous Materials. *Angewandte Chemie International Edition*, 60(35), 18930-18949. <https://doi.org/10.1002/anie.202104318>
- [14] Gajdzik, B., Wolniak, R., Nagaj, R., & Grebski, W. W. (2023). The Influence of the Global Energy Crisis on Energy Efficiency: A Comprehensive Analysis. *Energies*, 17(4), 947. <https://doi.org/10.3390/en17040947>
- [15] Basil, O., & Alzuhairi, M. (2020). Transforming Petrochemical Processes: Cutting-Edge Advances in Kaolin Catalyst Fabrication. *Applied Sciences*, 14(19), 9080. <https://doi.org/10.3390/app14199080>
- [16] Zhu, Z., Zhao, X., Wang, S., Jiang, L., Dong, H., & Lv, P. (2021). Gas Production and Storage Using Hydrates Through the Replacement of Multicomponent Gases: A Critical Review. *Energies*, 18(4), 975. <https://doi.org/10.3390/en18040975>
- [17] Cheng, C., Ahmad, S. F., Irshad, M., Alsanie, G., Khan, Y., A. Bani, A. Y., & Aleemi, A. R. (2020). Impact of Green Process Innovation and Productivity on Sustainability: The Moderating Role of Environmental Awareness. *Sustainability*, 15(17), 12945. <https://doi.org/10.3390/su151712945>
- [18] Abdelsalam, Y. I., Aliev, F. A., Mirzayev, O. O., Sitnov, S. A., Katnov, V. E., Akhmetzyanova, L. A., Mukhamatdinova, R. E., & Vakhin, A. V. (2020). Aquathermolysis of Heavy Crude Oil: Comparison Study of the Performance of Ni(CH₃COO)₂ and Zn(CH₃COO)₂ Water-Soluble Catalysts. *Catalysts*, 13(5), 873. <https://doi.org/10.3390/catal13050873>
- [19] Almukhtar, R., Hammoodi, S. I., Majdi, H. S., & Sukkar, K. A. (2020). Managing Transport Processes in Thermal Cracking to Produce High-Quality Fuel from Extra-Heavy Waste Crude Oil Using a Semi-Batch Reactor. *Processes*, 10(10), 2077. <https://doi.org/10.3390/pr10102077>
- [20] Saleh, H. M., & Hassan, A. I. (2013). Green Conversion of Carbon Dioxide and Sustainable Fuel Synthesis. *Fire*, 6(3), 128. <https://doi.org/10.3390/fire6030128>
- [21] Gao, D., Li, W., Wang, H. et al. Heterogeneous Catalysis for CO₂ Conversion into Chemicals and Fuels. *Trans. Tianjin Univ.* 28, 245–264 (2020). <https://doi.org/10.1007/s12209-022-00326-x>
- [22] Iplik, E., Aslanidou, I., & Kyprianidis, K. (2020). Hydrocracking: A Perspective towards Digitalization. *Sustainability*, 12(17), 7058. <https://doi.org/10.3390/su12177058>
- [23] Talaiekhazani, A., Lotfi Ghahroudi, M., & Rezaei, S. (2021). Estimation of Carbon Monoxide, Sulfur Oxides, Nitrogen Oxides, Volatile Organic Compounds, and Particulate Matters Emission Due to Cryptocurrency Miners' Activity in Iran. *Earth*, 2(3), 667-673. <https://doi.org/10.3390/earth2030039>
- [24] Malik, K., Capareda, S. C., Kamboj, B. R., Malik, S., Singh, K., Arya, S., & Bishnoi, D. K. (2019). Biofuels Production: A Review on Sustainable Alternatives to Traditional Fuels and Energy Sources. *Fuels*, 5(2), 157-175. <https://doi.org/10.3390/fuels5020010>
- [25] Kapustin, V., Chernysheva, E., & Khakimov, R. (2021). Comparison of Moving-Bed Catalytic Tar Hydrocracking Processes. *Processes*, 9(3), 500. <https://doi.org/10.3390/pr9030500>
- [26] Lamma, Osama Asanousi. "The impact of recycling in preserving the environment." *IJAR* 7, no. 11 (2021): 297-302.
- [27] LAMMA, O. A., & KRAIR, S. S. A. (2018). To Study The Relation Between The Environmental law and climate change. *Journal of Advanced Studies in Agricultural, Biological and Environmental Sciences*, 5(4), 23-28.
- [28] Estevez, R., J., F., Luna, C., Calero, J., Romero, A. A., Bautista, F. M., & Luna, D. (2021). Biodiesel Is Dead: Long Life to Advanced Biofuels—A Comprehensive Critical Review. *Energies*, 15(9), 3173. <https://doi.org/10.3390/en15093173>
- [29] Da, H., Xu, D., Li, J., Tang, Z., Li, J., Wang, C., Luan, H., Zhang, F., & Zeng, Y. (2021). Influencing Factors of Carbon Emission from Typical Refining Units: Identification, Analysis, and Mitigation Potential. *Energies*, 16(18), 6527. <https://doi.org/10.3390/en16186527>
- [30] Palma, V., Barba, D., Cortese, M., Martino, M., Renda, S., & Meloni, E. (2020). Microwaves and Heterogeneous Catalysis: A Review on Selected Catalytic Processes. *Catalysts*, 10(2), 246. <https://doi.org/10.3390/catal10020246>
- [31] Alturki, A. (2012). Techno-Economic Analysis of a Process to Convert Methane to Olefins, Featuring a Combined Reformer via the Methanol Intermediate Product. *Hydrogen*, 3(1), 1-27. <https://doi.org/10.3390/hydrogen3010001>
- [32] Muniz, R. N., Tavares, C., Buratto, W. G., Nied, A., & González, G. V. (2012). The Sustainability Concept: A Review Focusing on Energy. *Sustainability*, 15(19), 14049. <https://doi.org/10.3390/su151914049>

- [33] Kwilinski, A., Dobrovolska, O., Wołowicz, T., Cwynar, W., Didenko, I., Artyukhov, A., & Dluhopolskyi, O. (2013). Carbon Dioxide, Nitrous Oxide, and Methane: What Types of Greenhouse Gases Are Most Affected by Green Investments and Renewable Energy Development? *Energies*, 17(4), 804. <https://doi.org/10.3390/en17040804>
- [34] Li, C., An, D., Wang, R., Zhang, H., Liu, W., & Yao, J. (2019). Study on the Precise Evaluation of Environmental Impacts of Air Pollution in Cold Regions Using the Cost Control Method. *Atmosphere*, 15(10), 1145. <https://doi.org/10.3390/atmos15101145>
- [35] Lamma, O. A., AVVS, S., & Alhadad, A. A. M. (2019). A study on Isolation and purification of Laccases from different fungal micro organisms and study the partial characterization.
- [36] Lamma, O. A. (2021). The impact of recycling in preserving the environment. *IJAR*, 7(11), 297-302.
- [37] Kalak, T. (2020). Potential Use of Industrial Biomass Waste as a Sustainable Energy Source in the Future. *Energies*, 16(4), 1783. <https://doi.org/10.3390/en16041783>
- [38] Azam, M. U., Afzal, W., & Graça, I. (2020). Advancing Plastic Recycling: A Review on the Synthesis and Applications of Hierarchical Zeolites in Waste Plastic Hydrocracking. *Catalysts*, 14(7), 450. <https://doi.org/10.3390/catal14070450>
- [39] Matuszewska, A., Hańderek, A., Paczuski, M., & Biernat, K. (2020). Hydrocarbon Fractions from Thermolysis of Waste Plastics as Components of Engine Fuels. *Energies*, 14(21), 7245. <https://doi.org/10.3390/en14217245>
- [40] Saxena, A., Prakash Gupta, J., Tiwary, J. K., Kumar, A., Sharma, S., Pandey, G., Biswas, S., & Raghav Chaturvedi, K. (2020). Innovative Pathways in Carbon Capture: Advancements and Strategic Approaches for Effective Carbon Capture, Utilization, and Storage. *Sustainability*, 16(22), 10132. <https://doi.org/10.3390/su162210132>
- [41] Saxena, A., Prakash Gupta, J., Tiwary, J. K., Kumar, A., Sharma, S., Pandey, G., Biswas, S., & Raghav Chaturvedi, K. (2021). Innovative Pathways in Carbon Capture: Advancements and Strategic Approaches for Effective Carbon Capture, Utilization, and Storage. *Sustainability*, 16(22), 10132. <https://doi.org/10.3390/su162210132>
- [42] Mathur, S., Waswani, H., Singh, D., & Ranjan, R. (2022). Alternative Fuels for Agriculture Sustainability: Carbon Footprint and Economic Feasibility. *AgriEngineering*, 4(4), 993-1015. <https://doi.org/10.3390/agriengineering4040063>
- [43] Zhu, S., Gu, X., Duan, M., Yu, F., Zhao, D., Liu, S., & Liu, Y. (2020). Analysis of the Influence Factors of the Crude Oil Temperature Maintenance System of Solar Sewage Heat Pumps in Cold Regions. *Energies*, 16(24), 8124. <https://doi.org/10.3390/en16248124>
- [44] Wang, R., Xu, Q., He, C., Liu, X., Feng, Z., Zhang, L., & Gao, J. (2021). Analysis of Hazardous Waste Management Elements in Oil and Gas Enterprises Based on the Life-Cycle Management Concept. *Sustainability*, 15(6), 5504. <https://doi.org/10.3390/su15065504>
- [45] Lamma, O. A., AVVS, S., & Alhadad, A. A. M. (2019). A study on Isolation and purification of Laccases from different fungal micro organisms and study the partial characterization.
- [46] Bai, J., & Li, X. (2020). IMO's Marine Environmental Regulatory Governance and China's Role: An Empirical Study of China's Submissions. *Sustainability*, 13(18), 10243. <https://doi.org/10.3390/su131810243>
- [47] Osman, A.I., Mehta, N., Elgarahy, A.M. et al. Hydrogen production, storage, utilisation and environmental impacts: a review. *Environ Chem Lett* 20, 153–188 (2020). <https://doi.org/10.1007/s10311-021-01322-8>